

ASYMPTOTICS OF CHARACTERS OF SYMMETRIC GROUPS RELATED TO STANLEY-FÉRAY CHARACTER FORMULA

VALENTIN FÉRAY AND PIOTR ŚNIADY

ABSTRACT. We prove an upper bound for characters of the symmetric groups. Namely, we show that there exists a constant $a > 0$ with a property that for every Young diagram λ with n boxes, $r(\lambda)$ rows and $c(\lambda)$ columns

$$|\chi^\lambda(\pi)| \leq \left[a \max \left(\frac{r(\lambda)}{n}, \frac{c(\lambda)}{n}, \frac{|\pi|}{n} \right) \right]^{|\pi|},$$

where $|\pi|$ is the minimal number of factors needed to write $\pi \in S_n$ as a product of transpositions and

$$\chi^\lambda(\pi) = \frac{\text{Tr } \rho^\lambda(\pi)}{\text{Tr } \rho^\lambda(e)}$$

is the character of the symmetric group. We also give uniform estimates for the error term in the Vershik-Kerov's and Biane's character formulas and give a new formula for free cumulants of the transition measure.

1. INTRODUCTION

1.1. Normalized characters. For a Young diagram λ having n boxes and a permutation $\pi \in S_l$ (where $l \leq n$) we define the *normalized character*

$$(1) \quad \Sigma^\lambda(\pi) = (n)_l \chi^\lambda(\pi),$$

where $(n)_l = n(n-1) \cdots (n-l+1)$ denotes the falling power and where

$$\chi^\lambda(\pi) = \frac{\text{Tr } \rho^\lambda(\pi)}{\text{Tr } \rho^\lambda(e)}$$

is the character rescaled in such a way that $\chi^\lambda(e) = 1$.

1.2. Short history of the problem. Unfortunately, the canonical tool for calculating characters, the Murnaghan–Nakayama rule, quickly becomes cumbersome and hence intractable for computing characters corresponding to large Young diagrams. Nevertheless Roichman [Roi96] showed that it is possible to use it to find an upper bound for characters, namely he proved that there exist constants $0 < q < 1$ and $b > 0$ such that

$$(2) \quad |\chi^\lambda(\pi)| \leq \left[\max \left(\frac{r(\lambda)}{n}, \frac{c(\lambda)}{n}, q \right) \right]^{b |\text{supp } \pi|},$$

1

where $r(\lambda)$, $c(\lambda)$ denote the number of rows and columns of λ and $\text{supp } \pi$ denotes the support of a permutation π (the set of its non-fixed points). Inequality (2) is not satisfactory for many practical purposes (such as [MR06]) since it provides rather weak estimates in the case when the Young diagram λ is *balanced*, i.e. $r(\lambda), c(\lambda) = O(\sqrt{n})$ and π is not a very long permutation, i.e. $|\text{supp}(\pi)| = o(n)$.

Another approach to this problem was initiated by Biane [Bia98, Bia03] who showed that the value of the normalized character $\Sigma^\lambda(\pi)$ can be expressed as a relatively simple polynomial—called Kerov character polynomial—in *free cumulants* of the *transition measure* of a Young diagram λ . The work of Biane was based on previous contributions of Kerov [Ker93, Ker99] and Vershik. Since free cumulants of the transition measure have a nice geometric interpretation therefore Kerov polynomials are a perfect tool for study of the character $\chi^\lambda(\pi)$ in the limit when the permutation π is fixed and the Young diagram λ tends in some sense to infinity.

Unfortunately, despite much progress in this field [Śni06a, GR05] our understanding of Kerov polynomials is still not satisfactory; in particular it is not clear how to use Kerov polynomials in order to obtain non-trivial estimates on the characters $\chi^\lambda(\pi)$ when the length $|\pi|$ of the permutation $\pi \in S_n$ is comparable with n .

In a recent work of one of us with Rattan [RŚ06] we took yet another approach: thanks to the generalized Frobenius formula we showed that the value of a normalized character of a given Young diagram λ can be bounded from above by the value of the normalized character of a rectangular Young diagram $p \times q$ for suitably chosen p, q . For such a rectangular Young diagram the value of the normalized character can be explicitly calculated thanks to the formula of Stanley [Sta04] (which will be recalled as Theorem 4 in Section 2). In this way we proved that for each C there exists a constant D with a property that if $r(\lambda), c(\lambda) < C\sqrt{n}$ then

$$(3) \quad |\chi^\lambda(\pi)| < \left(\frac{D \max(1, \frac{|\pi|^2}{n})}{\sqrt{n}} \right)^{|\pi|},$$

where $|\pi|$ denotes the minimal number of factors necessary to write π as a product of transpositions. Inequality (3) gives a much better estimate than (2) for balanced Young diagrams and quite short permutation ($|\pi| = o(\sqrt{n})$) but it gives non-trivial estimates only if $\max(r(\lambda), c(\lambda)) < O(n^{3/4})$.

For the sake of completeness we point out that the studies of Kerov polynomials and of the Stanley character formula are very much related to each other [Bia03].

1.3. The main result. Our main result is the following inequality.

Theorem 1. *There exists a constant $a > 0$ with a property that for every Young diagram λ*

$$|\chi^\lambda(\pi)| \leq \left[a \max \left(\frac{r(\lambda)}{n}, \frac{c(\lambda)}{n}, \frac{|\pi|}{n} \right) \right]^{|\pi|},$$

where n denotes the number of boxes of λ .

It is easy to check that (3) is a consequence of this theorem and that it gives better estimates than (2) if $\frac{r(\lambda)}{n}, \frac{c(\lambda)}{n}, \frac{|\pi|}{n}$ are smaller than some positive constant. It is natural to ask what is the optimal value of the constant a . Asymptotics of characters of symmetric groups related to Thoma characters shows that $a \geq 1$.

1.4. Young diagrams. In the following we shall identify a Young diagram λ with the set of its boxes which we regard as a subset of $\mathbb{N}^2$ given by a graphical representation of λ according to the French notation; namely, for a partition $\lambda = (\lambda_1, \dots, \lambda_k)$ it is the set

$$(4) \quad \lambda = \bigcup_{1 \leq i \leq k} \{1, 2, \dots, \lambda_i\} \times \{i\} = \{(p, q) \in \mathbb{N}^2 : 1 \leq p \leq \lambda_q\}.$$

1.5. The main tool: reformulation of Stanley-Féray character formula.

Our main tool in our investigations will be the following reformulation of Stanley-Féray character formula.

The set of cycles of a permutation π is denoted by $C(\pi)$. For given permutations $\sigma_1, \sigma_2 \in S_l$ we shall consider colorings h of the cycles of σ_1 (where each cycle is colored by the number of some column of λ) and of the cycles of σ_2 (where each cycle is colored by the number of some row of λ). Formally, each such coloring can be viewed as a function $h : C(\sigma_1) \sqcup C(\sigma_2) \rightarrow \mathbb{N}$. We say that a coloring h is compatible with a Young diagram λ if for all $c_1 \in C(\sigma_1)$ and $c_2 \in C(\sigma_2)$ if $c_1 \cap c_2 \neq \emptyset$ then $(h(c_1), h(c_2)) \in \lambda$; in other words

$$(5) \quad 0 < h(c_1) \leq \lambda_{h(c_2)}$$

holds true for all $c_1 \in C(\sigma_1), c_2 \in C(\sigma_2)$ such that $c_1 \cap c_2 \neq \emptyset$.

Theorem 2 (The new formulation of Stanley-Féray character formula). *For any Young diagram λ and a permutation $\pi \in S_l$ (where $l \leq n$) the value of the normalized character (1) is given by*

$$(6) \quad \Sigma^\lambda(\pi) = \sum_{\substack{\sigma_1, \sigma_2 \in S_l, \\ \sigma_1 \sigma_2 = \pi}} (-1)^{|\sigma_1|} N^\lambda(\sigma_1, \sigma_2),$$




FIGURE 1. Bipartite graph associated to the factorization $(12) = (1)(2) \cdot (12)$.

where

$$(7) \quad N^\lambda(\sigma_1, \sigma_2) = \#\{h : h \text{ is a coloring of the cycles of } \sigma_1 \text{ and } \sigma_2 \text{ which is compatible with } \lambda\}.$$

Example 3. For a given factorization $\pi = \sigma_1 \sigma_2$ it is convenient to consider a bipartite graph with the set of vertices $C(\sigma_1) \sqcup C(\sigma_2)$ and with an edge between vertices $c_1 \in C(\sigma_1)$ and $c_2 \in C(\sigma_2)$ if and only if $c_1 \cap c_2 \neq \emptyset$. Notice that the value of $N^\lambda(\sigma_1, \sigma_2)$ does not depend on the exact form of σ_1 and σ_2 but only on the corresponding bipartite graph.

Figure 1 presents such a bipartite graph for $\pi = (12)$, $\sigma_1 = (1)(2)$, $\sigma_2 = (12)$. Now it becomes clear that

$$N^\lambda((1)(2), (12)) = \sum_i (\lambda_i)^2;$$

similarly

$$N^\lambda((12), (1)(2)) = \sum_i (\lambda'_i)^2,$$

where λ' denotes the Young diagram conjugate to λ . In this way Theorem 2 shows that

$$\begin{aligned} \Sigma^\lambda(12) &= n(n-1) \frac{\text{Tr } \rho^\lambda(12)}{\text{Tr } \rho^\lambda(e)} = \\ N^\lambda((1)(2), (12)) - N^\lambda((12), (1)(2)) &= \sum_i (\lambda_i)^2 - \sum_i (\lambda'_i)^2. \end{aligned}$$

1.6. Overview of the paper. In Section 2 we recall the original version of Stanley-Féray character formula. We also show that it is equivalent to Theorem 2 and we present its new proof.

In Section 3 we present a relation between the characters of symmetric groups and characters of some Gaussian random matrices. We also give a new formula for calculating free cumulants of (the transition measure of) a Young diagram.

Section 4 is devoted to the proofs of some technical inequalities.

In Section 5 we prove estimates for the characters of the symmetric groups based on Stanley-Féray character formula.

2. STANLEY-FÉRAY CHARACTER FORMULA

2.1. The original version of Stanley-Féray character formula. The direction of research presented in this paper was initiated by Stanley [Sta04] who proved the following result.

Theorem 4. *For any integers $p, q \geq 1$ the normalized character of the rectangular Young diagram $p \times q = \underbrace{(q, \dots, q)}_{p \text{ times}}$ is given on $\pi \in S_l$ by*

$$\Sigma^{p \times q}(\pi) = \sum_{\substack{\sigma_1, \sigma_2 \in S_l, \\ \sigma_1 \sigma_2 = \pi}} (-1)^{|\sigma_1|} q^{|C(\sigma_1)|} p^{|C(\sigma_2)|}.$$

Let $\mathbf{p} = (p_1, \dots, p_r)$ and $\mathbf{q} = (q_1, \dots, q_r)$ with $q_1 \geq \dots \geq q_r$ be two sequences of positive integers. We denote by $\mathbf{p} \times \mathbf{q}$ the multi-rectangular partition

$$(8) \quad \mathbf{p} \times \mathbf{q} = (\underbrace{q_1, \dots, q_1}_{p_1 \text{ times}}, \underbrace{q_2, \dots, q_2}_{p_2 \text{ times}}, \dots).$$

We denote by $S(l)^{(r)}$ the set of *colored permutations*, i.e. pairs (σ, ϕ) , where $\sigma \in S_l$ and $\phi : \{1, \dots, l\} \rightarrow \{1, \dots, r\}$ is a function constant on each cycle of σ . Alternatively, the coloring ϕ can be regarded as a function on $C(\sigma)$, the set cycles of σ . Given a colored permutation $(\sigma, \phi) \in S(l)^{(r)}$ and a non-colored one $\pi \in S_l$ we define their product $(\sigma, \phi) \cdot \pi = (\sigma\pi, \psi)$, where the coloring ψ is defined by

$$(9) \quad \psi(c) = \max_{a \in c} \phi(a),$$

where $c \in C(\sigma\pi)$.

The following generalization of Theorem 4 to multi-rectangular Young diagrams was conjectured by Stanley [Sta06] and proved by Féray [Fér06] and therefore we refer to it as Stanley-Féray character formula.

Theorem 5 (The original formulation of Stanley-Féray character formula). *The value of the normalized character is given on $\pi \in S_l$ by*

$$(10) \quad \Sigma^{\mathbf{p} \times \mathbf{q}}(\pi) = (-1)^l \sum_{(\sigma, \phi) \in S(l)^{(r)}} \left[\prod_{b \in C(\sigma)} p_{\phi(b)} \prod_{c \in C(\sigma\pi)} -q_{\psi(c)} \right],$$

where ψ was defined in (9).

2.2. Equivalence of the two formulations of Stanley-Féray character formula. In this section we will show that Theorem 2 implies Theorem 5 (the proof of the opposite implication is analogous and we leave it to the Reader).

Theorem 2 implies Theorem 5. Let permutations $\sigma_1, \sigma_2 \in S_l$ such that $\sigma_1 \sigma_2 = \pi^{-1}$ be given. We denote $\sigma = \sigma_2$.

The sequence $(\underbrace{1, \dots, 1}_{p_1 \text{ times}}, \underbrace{2, \dots, 2}_{p_2 \text{ times}}, \dots)$ can be viewed as a coloring of the set $\{1, \dots, p_1 + \dots + p_r\}$ with the colors $\{1, \dots, r\}$. In this way a function $h : C(\sigma_2) \rightarrow \{1, 2, \dots, p_1 + \dots + p_r\}$ defines a coloring ϕ of the permutation σ_2 . Clearly, there are $\prod_{b \in C(\sigma)} p_{\phi(b)}$ functions h which correspond to a given coloring ϕ .

We shall count now in how many ways a function $h : C(\sigma_2) \rightarrow \{1, 2, \dots, p_1 + \dots + p_r\}$ can be extended to a function $h : C(\sigma_1) \sqcup C(\sigma_2) \rightarrow \mathbb{N}$ compatible with σ_1, σ_2 . Condition (5) takes the form

$$\text{for all } c_1 \in C(\sigma_1), c_2 \in C(\sigma_2) \text{ if } c_1 \cap c_2 \neq \emptyset \text{ then } 0 < h(c_1) \leq q_{\phi(c_2)};$$

in other words for every $c_1 \in C(\sigma_1)$

$$0 < h(c_1) \leq \min_{\substack{c_2 \in C(\sigma_2), \\ c_1 \cap c_2 \neq \emptyset}} q_{\phi(c_2)} = \min_{a \in c_1} q_{\phi(a)} = q_{\psi(c_1)}$$

therefore the number of such extensions is equal to $\prod_{c \in C(\sigma\pi)} q_{\psi(c)}$.

In this way we proved that

$$\Sigma^{\mathbf{p} \times \mathbf{q}}(\pi) = \Sigma^{\mathbf{p} \times \mathbf{q}}(\pi^{-1}) = (-1)^l \sum_{(\sigma, \phi) \in S(l)^{(r)}} \left[\prod_{b \in C(\sigma)} p_{\phi(b)} \prod_{c \in C(\sigma\pi)} -q_{\psi(c)} \right].$$

□

2.3. New proof of the Stanley-Féray formula. Let λ be a Young diagram consisting of n boxes. In the following we will distinguish the symmetric group S_n which permutes the elements $\{1, \dots, n\}$ and the symmetric group $\tilde{S}_n$ which permutes the boxes of λ .

For a box $\square \in \lambda$ we denote by $r(\square) \in \mathbb{N}$ (respectively, $c(\square) \in \mathbb{N}$) the row (respectively, the column) of $\square$; in this way $\square = (c(\square), r(\square))$.

If $\sigma \in \tilde{S}_n$ has a property that if boxes $\square_1, \square_2$ are in the same row then $\sigma(\square_1), \sigma(\square_2)$ are not in the same column then we define the number of column inversions $\text{cinv}(\sigma)$ as the number of pairs $\square_1, \square_2$ such that $\sigma(\square_1), \sigma(\square_2)$ are in the same column, $r(\square_1) < r(\square_2)$ and $r(\sigma(\square_1)) > r(\sigma(\square_2))$. For $\sigma \in \tilde{S}_n$ which do not have this property we define $(-1)^{\text{cinv}(\sigma)} = 0$.

The following theorem gives a very esthetically appealing formula for the characters of the symmetric groups.

Theorem 6. *Let a Young diagram λ having n boxes and $\pi \in S_n$ be given. Let $\hat{\pi} \in \tilde{S}_n$ be a random permutation distributed with the uniform distribution on the conjugacy class defined by π . Then*

$$\chi^\lambda(\pi) = \mathbb{E}[(-1)^{\text{cinv}(\hat{\pi})}].$$

Proof. We denote

$$P_\lambda = \{\sigma \in \tilde{S}_n : \sigma \text{ preserves each row of } \lambda\},$$

$$Q_\lambda = \{\sigma \in \tilde{S}_n : \sigma \text{ preserves each column of } \lambda\}$$

and define

$$a_\lambda = \sum_{\sigma \in P_\lambda} \sigma \in \mathbb{C}[\tilde{S}_n],$$

$$b_\lambda = \sum_{\sigma \in Q_\lambda} (-1)^{|\sigma|} \sigma \in \mathbb{C}[\tilde{S}_n],$$

$$c_\lambda = b_\lambda a_\lambda.$$

It is well-known that $p_\lambda = \alpha_\lambda c_\lambda$ is an idempotent for a constant α_λ which will be specified later. Its image $V_\lambda = \mathbb{C}[\tilde{S}_n]p_\lambda$ under multiplication from the right on the regular representation gives a representation ρ^λ (where the symmetric group acts by left multiplication) associated to a Young diagram λ . It turns out that $\alpha_\lambda = \frac{\dim V_\lambda}{n!}$. It follows that for $\tilde{\pi} \in \tilde{S}_n$

$$(11) \quad n! \frac{\text{Tr } \rho^\lambda(\tilde{\pi})}{\dim V_\lambda} = \frac{\text{Tr } \rho^\lambda(\tilde{\pi}^{-1})}{\alpha_\lambda} = \frac{1}{\alpha_\lambda} \sum_{\mu \in \tilde{S}_n} \langle \delta_\mu, \tilde{\pi}^{-1} \delta_\mu p_\lambda \rangle =$$

$$\sum_{\mu \in \tilde{S}_n} \sum_{\tilde{\sigma}_1 \in Q_\lambda} \sum_{\tilde{\sigma}_2 \in P_\lambda} (-1)^{|\tilde{\sigma}_1|} [\mu = \tilde{\pi}^{-1} \mu \tilde{\sigma}_1 \tilde{\sigma}_2].$$

We define $\hat{\pi} = \mu^{-1} \tilde{\pi} \mu$. For such a permutation $\hat{\pi}$ there exists at most one factorization $\hat{\pi} = \tilde{\sigma}_1 \tilde{\sigma}_2$; by Young lemma this factorization exists if and only if $(-1)^{\text{cinv}(\hat{\pi})} \neq 0$. Furthermore, if such a factorization exists then $(-1)^{|\sigma_1|} = (-1)^{\text{cinv}(\sigma)}$. It follows that

$$n! \frac{\text{Tr } \rho^\lambda(\tilde{\pi})}{\dim V_\lambda} = \sum_{\mu \in \tilde{S}_n} (-1)^{\text{cinv}(\mu^{-1} \tilde{\pi} \mu)}.$$

As μ runs over all permutations, $\hat{\pi} = \mu^{-1} \tilde{\pi} \mu$ runs over all elements of the conjugacy class defined by π which finishes the proof. $\square$

For permutations $\sigma_1, \sigma_2 \in S_l$ we define $\tilde{N}^\lambda(\sigma_1, \sigma_2)$ as the number of one-to-one functions f from $\{1, \dots, l\}$ to the set of boxes of λ with a property that $r \circ f$ is constant on each cycle of σ_2 and $c \circ f$ is constant on each cycle of σ_1 .

Proposition 7. *Let λ be a Young diagram having n boxes. For any permutation $\pi \in S_l$ (where $l \leq n$)*

$$\Sigma^\lambda(\pi) = \sum_{\substack{\sigma_1, \sigma_2 \in S_l \\ \sigma_1 \sigma_2 = \pi}} (-1)^{|\sigma_1|} \tilde{N}^\lambda(\sigma_1, \sigma_2).$$

Proof. Let us consider the case $l = n$. Let $\tilde{\pi} \in \tilde{S}_n$ be any permutation with the same cycle structure as $\pi \in S_n$. Our starting point is the analysis of (11). Notice that the multiset of the values of $\mu^{-1} \tilde{\pi} \mu$ (over $\mu \in \tilde{S}_n$) coincides with the multiset of the values of $f \circ \pi \circ f^{-1}$ (over bijections f). We define $\sigma_i = f^{-1} \circ \tilde{\sigma}_i \circ f$; then condition $\mu = \tilde{\pi}^{-1} \mu \tilde{\sigma}_1 \tilde{\sigma}_2$ is equivalent to $\pi = \sigma_1 \sigma_2$. It is easy to check that $\tilde{\sigma}_1 \in Q_\lambda$ if and only if $c \circ f$ is constant on each cycle of σ_1 and $\tilde{\sigma}_2 \in P_\lambda$ if and only if $r \circ f$ is constant on each cycle of σ_2 . Thus

$$(12) \quad n! \frac{\text{Tr } \rho^\lambda(\tilde{\pi})}{\dim V_\lambda} = \sum_{\substack{\sigma_1, \sigma_2 \in S_n, \\ \sigma_1 \sigma_2 = \pi}} (-1)^{|\sigma_1|} \tilde{N}^\lambda(\sigma_1, \sigma_2)$$

and the proof in the case when $l = n$ is finished.

For a permutation $\sigma \in S_n$ we denote by $\text{supp } \sigma \subseteq \{1, \dots, n\}$ the support of a permutation (the set of non-fixed points). We claim that a factorization $\pi = \sigma_1 \sigma_2$ has a non-zero contribution to (12) only if $\text{supp } \sigma_1, \text{supp } \sigma_2 \subseteq \text{supp } \pi$. Indeed, if $m \in \text{supp } \sigma_1 \setminus \text{supp } \pi = \text{supp } \sigma_2 \setminus \text{supp } \pi$ then for any bijection f at least one of the following conditions hold true: $r(f(m)) \neq r(f(\sigma_2(m)))$ (in this case $r \circ f$ is not constant on the cycles of σ_2) or $c(f(m)) \neq c(f(\sigma_1(\sigma_2(m))))$ (in this case $c(f(\sigma_1(\sigma_2(m)))) \neq c(f(\sigma_2(m)))$) hence $c \circ f$ is not constant on the cycles of σ_1 .

It follows that if $\pi \in S_l$ then we may restrict the sum in (12) to factorizations $\pi = \sigma_1 \sigma_2$, where $\sigma_1, \sigma_2 \in S_l$ which finishes the proof. $\square$

For permutations $\sigma_1, \sigma_2 \in S_l$ we define $\hat{N}^\lambda(\sigma_1, \sigma_2)$ as the number of all functions $f : \{1, \dots, l\} \rightarrow \lambda$ (with values in the set of boxes of λ) with a property that $r \circ f$ is constant on each cycle of σ_2 and $c \circ f$ is constant on each cycle of σ_1 .

Lemma 8. *For any Young diagram λ and a permutation $\pi \in S_l$*

$$(13) \quad \sum_{\substack{\sigma_1, \sigma_2 \in S_l \\ \sigma_1 \sigma_2 = \pi}} (-1)^{|\sigma_1|} \tilde{N}^\lambda(\sigma_1, \sigma_2) = \sum_{\substack{\sigma_1, \sigma_2 \in S_l \\ \sigma_1 \sigma_2 = \pi}} (-1)^{|\sigma_1|} \hat{N}^\lambda(\sigma_1, \sigma_2).$$

Proof. Let a function $f : \{1, \dots, l\} \rightarrow \lambda$ be given which is not a one-to-one function. It follows that there exists a transposition $\mu \in S_l$ with a property

that f is constant on the orbits of μ . Function f contributes to the sum

$$\sum_{\substack{\sigma_1, \sigma_2 \in S_l \\ \sigma_1 \sigma_2 = \pi,}} (-1)^{|\sigma_1|} \tilde{N}^\lambda(\sigma_1, \sigma_2)$$

with multiplicity

$$(14) \quad \sum_{\substack{\sigma_1, \sigma_2 \in S_l \\ \sigma_1 \sigma_2 = \pi}} (-1)^{|\sigma_1|},$$

where the sum runs over pairs (σ_1, σ_2) with a property that $\sigma_1 \sigma_2 = \pi$ and $c \circ f$ is constant on each cycle of σ_1 and $r \circ f$ is constant on each cycle of σ_2 .

Map $(\sigma_1, \sigma_2) \mapsto (\sigma'_1, \sigma'_2)$ with $\sigma'_1 = \sigma_1 \mu$, $\sigma'_2 = \mu \sigma_2$ is an involution of the pairs (σ_1, σ_2) which contribute to (14); the only less trivial condition which should be verified is that $c \circ f$ is constant on each cycle of σ'_1 but this is equivalent to $c \circ f$ being constant on each cycle of $\sigma'^{-1}_1 = \mu \sigma^{-1}_1$.

Since $(-1)^{|\sigma_1|} = (-1) \cdot (-1)^{|\sigma'_1|}$ therefore the contributions of the pairs (σ_1, σ_2) and (σ'_1, σ'_2) to (14) cancel. In this way we proved that (14) is equal to zero which finishes the proof. $\square$

Proof of Theorem 2. Proposition 7 and Lemma 8 show that

$$\Sigma^\lambda(\pi) = \sum_{\substack{\sigma_1, \sigma_2 \in S_l \\ \sigma_1 \sigma_2 = \pi}} (-1)^{|\sigma_1|} \hat{N}^\lambda(\sigma_1, \sigma_2).$$

Now it is enough to notice that $N^\lambda(\sigma_1, \sigma_2) = \hat{N}^\lambda(\sigma_1, \sigma_2)$; the desired bijection is defined as follows: if $m \in \{1, \dots, l\}$ fulfills $m \in c_1 \cap c_2$ for $c_i \in C(\sigma_i)$ we set $f(m) = (h(c_1), h(c_2))$. $\square$

2.4. Generalization to Young diagrams on $\mathbb{R}^2_+$. We may identify a Young diagram with a subset of $\mathbb{R}^2$ given by a graphical representation of λ (according to the French notation). For example, for a partition $\lambda = (\lambda_1, \dots, \lambda_k)$ it is the set

$$(15) \quad \bigcup_{1 \leq i \leq k} [0, \lambda_i] \times [i-1, i].$$

In this way we may consider colorings h of the cycles of permutations σ_1 and σ_2 which instead of natural take real values. If we fix some numbering of the cycles in $C = C(\sigma_1) \sqcup C(\sigma_2)$ then any such coloring $h: C \rightarrow \mathbb{R}_+$ can be identified with an element of $\mathbb{R}_+^{|C|}$.

We define

$$(16) \quad N^\lambda(\sigma_1, \sigma_2) = \text{vol}\{h \in \mathbb{R}_+^{|C|} : h \text{ compatible with } \lambda\}.$$




FIGURE 2. Example of a skew Young diagram for which Theorem 2 is false.

Notice that in the case when $\lambda \subset \mathbb{R}^2$ is as prescribed by (15), the set of functions $h \in \mathbb{R}_+^{|C|}$ compatible with λ is a polyhedron hence there is no difficulty in defining its volume; furthermore definitions (7) and (16) give the same value.

The advantage of the definition (16) is that it allows to define characters for any bounded set $\lambda \in \mathbb{R}_+^2$, in particular for multi-rectangular Young diagrams $\mathbf{p} \times \mathbf{q}$ for general sequences $p_1, \dots, p_k \geq 0, q_1, \dots, q_k \geq 0$ which do not have to be natural numbers—just like in the original papers [Sta04, Sta06, Fér06].

It is very natural therefore to ask if Theorem 2 holds true also for skew Young diagrams. Unfortunately, this is not the case as it can be seen for the skew Young diagram λ from Figure 2 since

$$\Sigma^\lambda(12) = 12 \frac{\text{Tr } \rho^\lambda(12)}{\text{Tr } \rho^\lambda(e)} = \frac{12}{5}$$

which is not even an integer!

3. CHARACTERS OF SYMMETRIC GROUPS, RANDOM MATRICES AND FREE PROBABILITY

3.1. Stanley-Féray character formula and random matrices. For a Young diagram λ we consider a random matrix $T_\lambda = (t_{ij})$ such that

- its entries (t_{ij}) are independent random variables;
- if $(i, j) \in \lambda$ then t_{ij} is a complex centered Gaussian variable, that is to say that

$$\begin{aligned} \mathbb{E}(t_{ij}) &= 0, \\ \mathbb{E}(t_{ij} \overline{t_{ij}}) &= 1, \\ \mathbb{E}(t_{ij}^2) &= 0; \end{aligned}$$

- otherwise, if $(i, j) \notin \lambda$ then $t_{ij} = 0$.

Since T_λ has only finitely many non-zero entries we may identify it with its truncation $T_\lambda = (t_{ij})_{1 \leq i, j \leq N}$, where $N \geq r(\lambda), c(\lambda)$.

We are interested in this matrix because the moments of $T_\lambda T_\lambda^*$ are given by a formula which is very similar to the Stanley-Féray formula for characters (Theorem 2):

Theorem 9. *With the definitions above and $\pi \in S_l$ with a cycle decomposition $k_1, \dots, k_r$*

$$(17) \quad \mathbb{E}(\text{Tr}(T_\lambda T_\lambda^*)^{k_1} \dots \text{Tr}(T_\lambda T_\lambda^*)^{k_r}) = \sum_{\substack{\sigma_1, \sigma_2 \in S_l \\ \sigma_1 \sigma_2 = \pi}} N^\lambda(\sigma_1, \sigma_2).$$

Proof. The first step is to expand the product and the trace on the left-hand side:

$$\begin{aligned} & \mathbb{E}(\text{Tr}(T_\lambda T_\lambda^*)^{k_1} \dots \text{Tr}(T_\lambda T_\lambda^*)^{k_r}) \\ &= \mathbb{E} \left[\left(\sum_{i_1^1, j_1^1, \dots, i_{k_1}^1, j_{k_1}^1} t_{i_1^1 j_1^1} \overline{t_{i_2^1 j_1^1}} \dots t_{i_{k_1}^1 j_{k_1}^1} \overline{t_{i_1^1 j_{k_1}^1}} \right) \right. \\ & \quad \left. \dots \left(\sum_{i_1^r, j_1^r, \dots, i_{k_r}^r, j_{k_r}^r} t_{i_1^r j_1^r} \overline{t_{i_2^r j_1^r}} \dots t_{i_{k_r}^r j_{k_r}^r} \overline{t_{i_1^r j_{k_r}^r}} \right) \right] \\ &= \sum_{i_1, j_1, \dots, i_l, j_l} \mathbb{E} \left[\prod_{m=1}^l t_{i_m j_m} \overline{t_{i_{\pi(m)} j_m}} \right]. \end{aligned}$$

Since random variables (t_{ij}) are Gaussian we can apply Wick formula [Zvo97] to each summand; in order to do this we need to consider all ways of pairing factors (t_{i_m, j_m}) with factors $(\overline{t_{i_{\pi(m)} j_m}})$. Each such a pairing can be identified with a permutation $\sigma \in S_l$ therefore

$$\begin{aligned} \mathbb{E} \left[\prod_{m=1}^l t_{i_m j_m} \overline{t_{i_{\pi(m)} j_m}} \right] &= \sum_{\sigma \in S_l} \left[\prod_{m=1}^l \mathbb{E} \left(t_{i_{\sigma(m)} j_{\sigma(m)}} \overline{t_{i_{\pi(m)} j_m}} \right) \right] = \\ & \sum_{\sigma \in S_l} \prod_{m=1}^l [i_{\sigma(m)} = i_{\pi(m)}] [j_{\sigma(m)} = j_m] [(i_{\sigma(m)}, j_{\sigma(m)}) \in \lambda]. \end{aligned}$$

If we plug this in our calculation,

$$\begin{aligned} & \mathbb{E}(\text{Tr}(T_\lambda T_\lambda^*)^{k_1} \dots \text{Tr}(T_\lambda T_\lambda^*)^{k_r}) = \\ & \sum_{\sigma \in S_l} \sum_{i_1, j_1, \dots, i_l, j_l} \left(\prod_{m=1}^l [i_{\sigma(m)} = i_{\pi(m)}] [j_{\sigma(m)} = j_m] [(i_{\sigma(m)}, j_{\sigma(m)}) \in \lambda] \right). \end{aligned}$$

If we denote $\sigma_1 = \pi \sigma_2^{-1}$, $\sigma_2 = \sigma$ then the sum over σ can be seen as a sum over all $\sigma_1, \sigma_2 \in S_l$ such that $\sigma_1 \sigma_2 = \pi$. If a sequence $i_1, \dots, i_l$ (respectively, sequence $j_1, \dots, j_l$) contributes to the above sum then it must be constant on each cycle of σ_1 (respectively, each cycle of σ_2). It follows that there is a bijective correspondence between sequences $i_1, j_1, \dots, i_l, j_l$

which contribute to the above sum and colorings of the cycles of σ_1 and σ_2 which are compatible with λ . $\square$

By comparing the above result with Theorem 2 we obtain the following corollary.

Corollary 10. *Let λ be a Young diagram. Then for any permutation $\pi \in S_l$ with a cycle decomposition $k_1, \dots, k_r$*

$$|\Sigma^\lambda(\pi)| \leq \mathbb{E}(\mathrm{Tr}(T_\lambda T_\lambda^*)^{k_1} \dots \mathrm{Tr}(T_\lambda T_\lambda^*)^{k_r}).$$

We shall not follow this idea in this article and we will prove all estimates from scratch, but it is worth noticing that the above Corollary shows that the asymptotics of characters of symmetric groups can be deduced from the corresponding asymptotics of random matrices. In particular it follows that when the lengths $k_1, \dots, k_r$ of the cycles of π are big enough then the asymptotics of the corresponding character is related to the limit distribution of the largest eigenvalue of $T_\lambda T_\lambda^*$ [SS98]. Notice, however, that due to the minus sign in Theorem 2 and the resulting cancelations the character $|\Sigma^\lambda(\pi)|$ could, at least in principle, be much smaller than the appropriate moment of the random matrix $T_\lambda T_\lambda^*$.

3.2. Formal Young diagrams. In the same way as Theorem 2 implies Theorem 5 equation (17) implies the following result.

Corollary 11. *If $\lambda = \mathbf{p} \times \mathbf{q}$ is a multirectangular Young diagram (8) then (18)*

$$\mathbb{E}(\mathrm{Tr}(T_\lambda T_\lambda^*)^{k_1} \dots \mathrm{Tr}(T_\lambda T_\lambda^*)^{k_r}) = \sum_{(\sigma, \phi) \in S(l)^{(r)}} \left[\prod_{b \in C(\sigma)} p_{\phi(b)} \prod_{c \in C(\sigma\pi)} q_{\psi(c)} \right].$$

It is somewhat disturbing that both the right-hand side of the original Stanley-Féray formula (10) and the right-hand side of (18) are polynomials $\mathbf{p}$ and $\mathbf{q}$ and as such make sense for arbitrary values of $\mathbf{p}$ and $\mathbf{q}$, in particular for negative values. If we allow use of such *formal Young diagrams* $\mathbf{p} \times \mathbf{q}$ (the term *generalized Young diagram* is already in use and it means *continuous Young diagram*) then the following result follows.

Theorem 12. *For any formal Young diagram $\mathbf{p} \times \mathbf{q}$ and a permutation $\pi \in S_l$ with a cycle decomposition $k_1, \dots, k_r$*

$$\Sigma^{\mathbf{p} \times \mathbf{q}}(\pi) = (-1)^l \mathbb{E}(\mathrm{Tr}(T_{\mathbf{p} \times (-\mathbf{q})} T_{\mathbf{p} \times (-\mathbf{q})}^*)^{k_1} \dots \mathrm{Tr}(T_{\mathbf{p} \times (-\mathbf{q})} T_{\mathbf{p} \times (-\mathbf{q})}^*)^{k_r}).$$

The authors of this article are still confused about the meaning of the above equality since (except for the case of an empty Young diagram) at least one of the formal Young diagrams $\mathbf{p} \times \mathbf{q}$, $\mathbf{p} \times (-\mathbf{q})$ does not make sense as a Young diagram.

3.3. Free cumulants of the transition measure. For a (continuous) Young diagram λ we denote by μ^λ its transition measure (which is a probability measure on the real line) [Ker93, Bia98] and by $R_m^\lambda := R_m(\mu^\lambda)$ we denote the m -th free cumulant of μ^λ . The importance of free cumulants R_m^λ in the study of the asymptotics of symmetric groups was pointed out by Biane [Bia98]. The following theorem gives a new formula for the free cumulants R_m^λ . It has a big advantage that it does not involve the notion of the transition measure and it is related directly with the shape of a Young diagram.

Theorem 13. *For any Young diagram λ*

$$(19) \quad R_{l+1}^\lambda = \sum_{\substack{\sigma_1, \sigma_2 \in S_l, \\ \sigma_1 \sigma_2 = (1, 2, \dots, l), \\ |\sigma_1| + |\sigma_2| = |(1, 2, \dots, l)|}} (-1)^{|\sigma_1|} N^\lambda(\sigma_1, \sigma_2),$$

where the sum runs over minimal factorizations of a cycle of length l .

Proof. For a Young diagram λ and $c > 0$ we denote by $c\lambda$ the (generalized) Young diagram obtained from λ by similarity with scale c . A function f on the set of (generalized) Young diagrams is said to be homogeneous of degree m if

$$f(c\lambda) = c^m f(\lambda)$$

holds true for all choices of λ and c . Each free cumulant R_m is homogenous of degree m .

The value of the normalized character $\Sigma^\lambda(1, 2, \dots, l)$ on a cycle can be expressed as a polynomial (known as Kerov polynomial) in free cumulants $(R_m^\lambda : m \in \{2, 3, \dots\})$:

$$\Sigma^\lambda(1, 2, \dots, l) = R_{l+1}^\lambda + (\text{terms of lower degree})$$

therefore R_{l+1}^λ is the homogeneous part of $\Sigma^\lambda(1, 2, \dots, l)$ with degree $l + 1$. We apply (6) for $\pi = (1, 2, \dots, l)$; it is easy to see that each summand on the right-hand side is homogeneous of degree $|C(\sigma_1)| + |C(\sigma_2)|$ which finishes the proof. $\square$

3.4. Generalized circular operators. Let $\mathcal{D}$ be the algebra of continuous functions on $\mathbb{R}_+$. We equip it with an expected value $\phi : \mathcal{D} \rightarrow \mathbb{C}$ given by $\phi(f) = \int_0^\infty f(t) dt$.

We consider an operator-valued probability space, which by definition is some $*$ -algebra $\mathcal{A}$ such that $\mathcal{D} \subseteq \mathcal{A}$ and equipped with a conditional expectation $\mathbb{E} : \mathcal{A} \rightarrow \mathcal{D}$. For a given (generalized) Young diagram λ let $T \in$

$\mathcal{A}$ be a generalized circular operator [VDN92, Spe98] with a covariance

$$\begin{aligned}
 [k(T, fT^*)](s) &= \int_{t:(t,s) \in \lambda} f(t) dt, \\
 [k(T^*, fT)](s) &= - \int_{t:(s,t) \in \lambda} f(t) dt, \\
 [k(T, fT)](s) &= 0, \\
 [k(T^*, fT^*)](s) &= 0.
 \end{aligned}
 \tag{20}$$

Theorem 14. *For any (generalized) Young diagram λ*

$$R_{l+1}^\lambda = \phi[(T^*T)^l].$$

Proof. It is easy to check that for permutations σ_1, σ_2 which contribute to (19) the corresponding bipartite graph is a tree therefore the calculation of $N^\lambda(\sigma_1, \sigma_2)$ is particularly simple, namely it is a certain iterated integral. The same iterated integral appears in the nested evaluation of amalgamated free cumulants therefore $N^\lambda(\sigma_1, \sigma_2) = \pm \phi[k_{\sigma_2}(T^*, T, \dots, T^*, T)]$. The plus/minus sign is due to the minus sign in the covariance (20). It is easy to check that in fact

$$(-1)^{|\sigma_1|} N^\lambda(\sigma_1, \sigma_2) = \phi[k_{\sigma_2}(T^*, T, \dots, T^*, T)].$$

The moment-cumulant formula

$$\mathbb{E}[(T^*T)^l] = \sum_{\sigma \in \text{NC}_2} k_\sigma(T^*, T, \dots, T^*, T)$$

finishes the proof since there is a bijective correspondence between non-crossing pair partitions and the minimal factorizations. $\square$

Except for the minus sign on the right-hand side of (20) the covariance of T looks in the same way as the covariance of generalized circular operators corresponding to the limits of Gaussian band matrices [Shl96] which should not be surprising in the light of Theorem 9. We refer also to the articles [Śni03, DH04] where a similar circular operator is considered.

Due to the analytic machinery of free probability the calculation of the moments of T in a closed form is possible in many cases therefore Theorem 14 gives a practical method of calculating the free cumulants of the Young diagrams.

4. TECHNICAL ESTIMATES

4.1. Estimates for the number of colorings $N^\lambda(\sigma_1, \sigma_2)$. As we already mentioned in Example 3 to permutations σ_1, σ_2 we can associate a bipartite graph $C(\sigma_1) \sqcup C(\sigma_2)$ with an edge between $c_1 \in C(\sigma_1)$ and $c_2 \in C(\sigma_2)$ if $c_1 \cap c_2 \neq \emptyset$.

For a bipartite graph $G = C_1 \sqcup C_2$ (not necessarily arising from the above construction) and a Young diagram λ we define $N^\lambda(G)$ as the number of colorings h of the vertices of $C_1 \sqcup C_2$ which are compatible with the Young diagram λ (the definition of compatibility in this context is a natural extension of the old one, i.e. we require that if $c_1 \in C_1$ and $c_2 \in C_2$ are connected by an edge then $(h(c_1), h(c_2)) \in \lambda$).

We denote by $G_{p,q}$ a full bipartite graph for which $|C_1| = p$ and $|C_2| = q$.

Lemma 15. *Let G be a finite bipartite graph with a property that the degree of any vertex is non-zero. It is possible (not necessarily in a unique way) to remove some of the edges of G in such a way that the resulting graph $\tilde{G}$ is a disjoint union of the graphs of the form $G_{1,1}$, $G_{k,1}$, $G_{1,k}$.*

Assume that a Young diagram λ consists of n boxes. Then, for any $A \geq r(\lambda), c(\lambda)$

$$(21) \quad N^\lambda(G) \leq A^{(\text{number of vertices of } G)} \left(\frac{n}{A^2} \right)^{(\text{number of connected components of } \tilde{G})}.$$

Proof. If the graph G contains an edge which connects two vertices of degree bigger than one we remove it and iterate this procedure; if no such edge exists then the resulting graph $\tilde{G}$ has the desired property.

Clearly, $N^\lambda(G) \leq N^\lambda(\tilde{G})$ therefore it is enough to find a suitable upper bound for $N^\lambda(\tilde{G})$. Since both sides of (21) are multiplicative with respect to the disjoint sum of graphs it is enough to prove (21) for $\tilde{G} \in \{G_{1,1}, G_{k,1}, G_{1,k}\}$. It is easy to verify that

$$\begin{aligned} N^\lambda(G_{1,1}) &= n, \\ N^\lambda(G_{k,1}) &= \sum_i \lambda_i^k \leq A^{k-1}n, \\ N^\lambda(G_{1,k}) &= \sum_i \lambda_i^k \leq A^{k-1}n \end{aligned}$$

which finishes the proof. $\square$

Proposition 16. *Suppose that $r(\lambda), c(\lambda) \leq A \leq n$ and $\sigma_1, \sigma_2 \in S_l$ and $\pi = \sigma_1 \sigma_2$. Then*

$$(22) \quad N^\lambda(\sigma_1, \sigma_2) \leq A^{|C(\sigma_1)| + |C(\sigma_2)|} \left(\frac{n}{A^2} \right)^{\text{orbits}(\sigma_1, \sigma_2)} \leq A^{l - |C(\pi)|} n^{|C(\pi)|} \left(\frac{1}{A} \right)^{o(\sigma_1, \sigma_2)},$$

where $\text{orbits}(\sigma_1, \sigma_2)$ denotes the number of orbits in the action of $\langle \sigma_1, \sigma_2 \rangle$ on the set $\{1, \dots, n\}$ and

$$o(\sigma_1, \sigma_2) = l - |C(\sigma_1)| - |C(\sigma_2)| + \text{orbits}(\sigma_1, \sigma_2).$$

Proof. The first inequality is a simple corollary from Lemma 15 since $A^2 \geq r(\lambda)c(\lambda) \geq n$ and the number of connected components of $\tilde{G}$ is bounded from below by $\text{orbits}(\sigma_1, \sigma_2)$. The second inequality follows by multiplying by

$$\left(\frac{n}{A}\right)^{|C(\pi)| - \text{orbits}(\sigma_1, \sigma_2)} \geq 1.$$

□

4.2. Estimates for the number of factorizations. Now, we have to find a bound of the number of factorizations of π with a given value of the statistic $o(\sigma_1, \sigma_2)$.

Lemma 17. *Let $\pi, \sigma_1, \sigma_2 \in S_l$ be such that $\pi = \sigma_1\sigma_2$. There exist permutations $\sigma'_1, \sigma'_2 \in S_l$ such that $\pi = \sigma'_1\sigma'_2$, $|\sigma'_1| + |\sigma'_2| = |\sigma_1| + |\sigma_2|$, $|\sigma'_2| = |\sigma'_2\sigma_2^{-1}| + |\sigma_2|$ and every cycle of σ'_1 is contained in some cycle of σ'_2 . Furthermore, $|\sigma'_1| = o(\sigma_1, \sigma_2)$.*

Proof. If every cycle of σ_1 is contained in some cycle of σ_2 then $\sigma'_1 = \sigma_1$ and $\sigma'_2 = \sigma_2$ have the required property.

Otherwise, there exist $a, b \in \{1, \dots, n\}$ such that a and b belong to the same cycle of σ_1 but not to the same cycle of σ_2 . We define $\sigma'_1 = \sigma_1(a, b)$, $\sigma'_2 = (a, b)\sigma_2$. Notice that $|\sigma'_1| = |\sigma_1| - 1$, $|\sigma'_2| = |\sigma_2| + 1$, and the partitions Π and Π' of $\{1, \dots, l\}$ in the orbits under the action of the subgroups $\langle \sigma_1, \sigma_2 \rangle$ and $\langle \sigma'_1, \sigma'_2 \rangle$ are the same.

We iterate this procedure if necessary (it will finish after a finite number of steps because the length of σ_1 decreases in each step). It remains to prove that $|\sigma'_2| \geq |\sigma'_2\sigma_2^{-1}| + |\sigma_2|$ (the opposite inequality follows from the triangle inequality): notice that $|\sigma'_2| - |\sigma_2|$ is equal to k (where k is the number of steps after which the procedure has terminated) and $\sigma'_2\sigma_2^{-1}$ is a product of k transpositions, hence $|\sigma'_2\sigma_2^{-1}| \leq k$.

Furthermore, as every cycle of σ'_1 is contained in some cycle of σ'_2 , the partition Π' is just $C(\sigma'_2)$, so $|C(\sigma'_2)| = \text{orbits}(\sigma'_1, \sigma'_2)$ therefore

$$o(\sigma_1, \sigma_2) = o(\sigma'_1, \sigma'_2) = l - |C(\sigma'_1)| = |\sigma'_1|.$$

□

Lemma 18. *For any integers $l \geq 1$ and $i \geq 0$ and for any $\pi \in S_l$*

$$(23) \quad \#\{\sigma \in S_l : |\sigma| = i\} \leq \frac{l^{2i}}{i!}.$$

Proof. Since every permutation in S_l appears exactly once in the product

$$[1 + (12)][1 + (13) + (23)] \cdots [1 + (1l) + \cdots + (l-1, l)],$$

we have

$$\sum_i x^i \# \{ \sigma \in S_l : |\sigma| = i \} = (1+x)(1+2x) \cdots (1+(l-1)x).$$

Each of the coefficients of x^k on the right-hand side is bounded from above by the corresponding coefficient of $e^x e^{2x} \cdots e^{(l-1)x} = e^{\frac{l(l-1)x}{2}}$, finishing the proof. $\square$

Lemma 19. *There exists a constant C_0 with a property that for any k the number of minimal factorisations $\sigma_1 \sigma_2 = \pi$, $|\sigma_1| + |\sigma_2| = |\pi|$ of a cycle $\pi = (1, \dots, k)$ and such that the associated graph $\tilde{G}$ consists of $s \geq 2$ components is bounded from above by*

$$\frac{(C_0 k)^{2s-2}}{(2s-2)!}.$$

Proof. Since the factorization is minimal therefore the graph G associated to σ_1, σ_2 is a tree.

In each connected component of $\tilde{G}$ there is at most one vertex of degree higher than one and we shall decorate this vertex. If in some connected component of $\tilde{G}$ there are no such vertices we decorate any of them. In this way the decorated vertices can be identified with connected components of $\tilde{G}$.

We will give to G a structure of planted planar tree: the root is the cycle of σ_1 containing 1 and his left-most edge links it to the cycle of σ_2 containing 1.

We consider the graph G' obtained from G by removing the leaves (except the root) and the graph G'' which consists of the decorated vertices of $\tilde{G}$; we connect two vertices $A, B \in G''$ by an edge if vertices A, B are connected in G (or, equivalently, G') by a *direct path*, i.e. a path which does not pass through any connected component of $\tilde{G}$ other than the ones specified by A and B . It is easy to see that G'' inherits the structure of a plane rooted tree from G (we define the root of G'' to be the connected component of $\tilde{G}$ of the root of G) and it has s vertices. It follows that the number of such trees G'' is bounded from above by the Catalan number $\frac{1}{s+1} \binom{2s}{s} < 4^s$.

In order to reconstruct tree G' from G'' we have to specify for each edge of G'' if it comes from a single edge of G' or from a pair or a triple of consecutive edges of G' ; it follows that we have (at most) 3^{s-1} choices. It might happen also for two adjacent (with respect to the planar structure) edges e_1, e_2 of G'' that each of these edges e_i corresponds to a pair or a triple of consecutive edges $f_i = (f_{i1}, f_{i2}, f_{i3})$ of G' and these tuples f_1 and f_2 have one edge in common. There are at most $2s-3$ such pairs of adjacent edges which accounts for at most 2^{2s-3} choices. If the root of G

is not a decorated vertex, it might happen that it is a leaf or that it belong to the left-most and/or to the right-most edge of the root of G' : there are 4 choices for that.

In order to reconstruct tree G from G' we have to specify if the root of G is a decorated vertex or not. Furthermore we have to specify places in which we will add missing l leaves to the tree G' (note that $l \leq k+1-s$); it is easy to see that this is equivalent to specifying a partition $l = a_1 + \dots + a_{2s-1}$, where $a_1, \dots, a_{2s-1} \geq 0$ are integers. It follows that the number of choices is bounded from above by

$$2 \binom{l+2s-2}{2s-2} \leq 2 \binom{k+s-1}{2s-2} \leq 2 \frac{k^{2s-2}}{(2s-2)!}.$$

A minimal factorisation is determined by its bicolored graph with a marked edge, for example the one linking the two cycles containing 1 [GJ92]. With our construction, the coloring is determined by the root which is always white (i.e. a cycle of σ_1) and the marked edge is its left-most edge.

It follows that the total number of choices is bounded from above by

$$2 \cdot 3^{s-1} \cdot 2^{2s-1} \cdot 4^s \frac{k^{2s-2}}{(2s-2)!}.$$

□

5. ASYMPTOTICS OF CHARACTERS

5.1. Upper bound for characters: proof of Theorem 1.

Proof of Theorem 1. Let $k_1, \dots, k_r \geq 2$ be the lengths of the non-trivial cycles in the cycle decomposition of $\pi \in S_n$. It follows that $l := k_1 + \dots + k_r = |\text{supp } \pi|$ and in the following we will regard π as an element of S_l . We denote $A = \max(l, r(\lambda), c(\lambda))$.

We consider a map which to a pair (σ_1, σ_2) associates any pair (σ'_1, σ'_2) as prescribed by Lemma 17. For any fixed σ'_2 the permutations σ_2 such that $|\sigma'_2| = |\sigma'_2 \sigma_2^{-1}| + |\sigma_2|$ can be identified with non-crossing partitions of the cycles of σ'_2 (see [Bia97, Section 1.3]). It follows that the number of such permutations σ_2 is equal to the product of appropriate Catalan numbers and, hence, this product is bounded from above by 4^l . Therefore Theorem 2 and

Proposition 16 show that

$$\begin{aligned} |\Sigma^\lambda(\pi)| &\leq \sum_{\substack{\sigma_1, \sigma_2 \in S_l, \\ \sigma_1 \sigma_2 = \pi}} A^{l-r} n^r \left(\frac{1}{A}\right)^{o(\sigma_1, \sigma_2)} \leq 4^l \sum_{\substack{\sigma'_1, \sigma'_2 \in S_l, \\ \sigma'_1 \sigma'_2 = \pi}} A^{l-r} n^r \left(\frac{1}{A}\right)^{|\sigma'_1|} \\ &\leq 4^l A^{l-r} n^r \sum_{\sigma'_1 \in S_l} \left(\frac{1}{A}\right)^{|\sigma'_1|} \leq 4^l A^{l-r} n^r \sum_{i \geq 0} \frac{l^{2i}}{A^i i!}, \end{aligned}$$

where the last inequality follows from Lemma 18. It follows that

$$(24) \quad |\Sigma^\lambda(\pi)| \leq 4^l A^{l-r} n^r e^{\frac{l^2}{A}} \leq (4e)^l A^{l-r} n^r.$$

After dividing by $(n)_l \geq \left(\frac{n}{e}\right)^l$ we obtain

$$(25) \quad |\chi^\lambda(\pi)| \leq (4e^2)^l \left(\frac{A}{n}\right)^{l-r}$$

which finishes the proof because $l = |\text{supp } \pi| \leq 2|\pi|$ and $l - r = |\pi|$. $\square$

5.2. Error term for balanced Young diagrams. Biane [Bia98] proved that if the permutation π is fixed and the Young diagram λ is balanced (i.e. $r(\lambda), c(\lambda) = O(\sqrt{n})$) then

$$\chi^\lambda(\pi) = \frac{1}{(n)_{|\text{supp}(\pi)|}} \prod_{i=1}^r R_{k_i+1}(\lambda) + O\left(n^{-\frac{|\pi|+2}{2}}\right).$$

The following theorem gives a uniform estimate for the error term in Biane's formula (note that if π is fixed and the Young diagram balanced, with $\epsilon = \frac{C}{\sqrt{n}}$ we recover the result of Biane).

Theorem 20. *There exists a constant a such that, for any $0 < \varepsilon < 1$, any Young diagram λ of size n and any permutation $\pi \in S_n$ such that $|\text{supp}(\pi)|^2 \leq \varepsilon A$ and $r(\lambda), c(\lambda) \leq A \leq n$ we have:*

$$\left| \chi^\lambda(\pi) - \frac{1}{(n)_{|\text{supp}(\pi)|}} \prod_{i=1}^r R_{k_i+1}(\lambda) \right| \leq \left(\varepsilon^2 + \frac{A}{n} \varepsilon \right) \left(\frac{aA}{n} \right)^{|\pi|},$$

where the k_i are the lengths of the non-trivial cycles of π .

Proof. Using Theorem 2 and Theorem 13 together with the fact that any minimal factorisation of π is a product of minimal factorisations of its cycles, we can write:

$$\Sigma^\lambda(\pi) - \prod_{i=1}^r R_{k_i+1}^\lambda = \sum_{\substack{\sigma_1, \sigma_2 \in S_l \\ \sigma_1 \sigma_2 = \pi \\ |\sigma_1| + |\sigma_2| > |\pi|}} (-1)^{|\sigma_1|} N^\lambda(\sigma_1, \sigma_2),$$

where $l = |\text{supp } \pi|$ is the support of π . To such a pair (σ_1, σ_2) of permutations we can associate one of the pairs of permutations (σ'_1, σ'_2) given by Lemma 17 with $|\sigma'_1| \geq 1$.

Consider the case $|\sigma'_1| = 1$. Then $\text{orbits}(\sigma_1, \sigma_2) = \text{orbits}(\sigma'_1, \sigma'_2) \geq |C(\pi)| - 1$ and the first inequality in (22) shows that

$$N^\lambda(\sigma_1, \sigma_2) \leq A^{|C(\sigma_1)|+|C(\sigma_2)|} \left(\frac{n}{A^2} \right)^{|C(\pi)|-1}$$

therefore the estimate given by Proposition 16 can be improved to the following one:

$$N^\lambda(\sigma_1, \sigma_2) \leq A^{l-|C(\pi)|} n^{|C(\pi)|} \frac{1}{n}.$$

Clearly $A \geq l$ therefore by the same argument as in the proof of Theorem 1, we obtain the inequality

$$\left| \Sigma^\lambda(\pi) - \prod_{i=1}^r R_{k_i+1}(\lambda) \right| \leq 4^l A^{l-r} n^r \left(\frac{l^2}{n} + \sum_{i \geq 2} \frac{l^{2i}}{A^i i!} \right).$$

The proof is now finished thanks to the remarks of the previous subsection and the inequality $\exp(z) - 1 - z \leq z^2$ for $0 < z < 1$. $\square$

5.3. Characters of symmetric groups related to Thoma characters. Vershik and Kerov [VK81] proved that if π is a fixed permutation with the lengths of non-trivial cycles $k_1, \dots, k_r$ then for any Young diagram λ with n boxes

$$\chi^\lambda(\pi) = \prod_{i=1}^r \left[\sum_j \alpha_j^{k_i} - \sum_j (-\beta_j)^{k_i} \right] + O\left(\frac{1}{n}\right)$$

where $\alpha_j = \frac{\lambda_j}{n}$, $\beta_j = \frac{\lambda'_j}{n}$; we prefer to write this formula in an equivalent form

$$(26) \quad \chi^\lambda(\pi) = \frac{n^l}{(n)_l} \prod_{i=1}^r \left[\sum_j \alpha_j^{k_i} - \sum_j (-\beta_j)^{k_i} \right] + O\left(\frac{1}{n}\right).$$

In this section we will prove Theorem 21 which together with Theorem 20 give a uniform estimate for the error term in the formula (26). In particular, for $A = n$ and $\epsilon = \frac{C}{n}$ we recover the result of Vershik and Kerov.

Theorem 21. *There exist constants $a, C > 0$ with the following property. Let $k_1, \dots, k_r$ be positive integers; we denote $k_1 + \dots + k_r = l$. If λ is a Young diagram having n boxes with less than A boxes in each row and each column and such that $\epsilon = \frac{(k_1^2 + \dots + k_r^2)n}{A^2} < C$ then*

$$(27) \quad \left| \frac{\prod_{i=1}^r R_{k_i+1}^\lambda}{n^l} - \prod_{i=1}^r \left[\sum_j \alpha_j^{k_i} - \sum_j (-\beta_j)^{k_i} \right] \right| \leq \epsilon \left(\frac{A}{n} \right)^{l-r} a^r,$$

where $\alpha_j = \frac{\lambda_j}{n}$, $\beta_i = \frac{\lambda'_i}{n}$.

Proof. Firstly, let us consider the case $r = 1$. Note that

$$N^\lambda(e, (1, \dots, k)) = \sum_j (n\alpha_j)^k,$$

$$N^\lambda((1, \dots, k), e) = \sum_j (n\beta_j)^k$$

therefore Theorem 13 implies that the left-hand side of (27) is equal to

$$\left| \frac{1}{n^k} \sum_{\substack{\sigma_1, \sigma_2 \in S_k \setminus \{e\} \\ \sigma_1 \sigma_2 = \pi \\ |\sigma_1| + |\sigma_2| = |\pi|}} (-1)^{|\sigma_1|} N^\lambda(\sigma_1, \sigma_2) \right|.$$

Let a pairs of permutations σ_1, σ_2 which contribute to the above sum be fixed; we consider the bipartite graph G and the graph $\tilde{G}$ given by Lemma 15. Clearly, in this case graph $\tilde{G}$ has more than one component. With Lemma 19 and Lemma 15

$$(28) \quad \left(\frac{n}{A} \right)^{k-1} \left| \frac{R_{k+1}^\lambda}{n^k} - \sum_j (\alpha_j^k - (-\beta_j)^k) \right| \leq$$

$$\sum_{s \geq 2} \frac{(C_0 k)^{2s-2}}{(2s-2)!} \left(\frac{n}{A^2} \right)^{s-1} \leq \frac{2C_0^2 k^2 n}{A^2} = 2C_0^2 \varepsilon,$$

where the last inequality holds true if $\frac{2C_0^2 k^2 n}{A^2} = 2C_0^2 \varepsilon$ is smaller than some positive constant and the proof is finished in the case $r = 1$.

For the general case, we put $\epsilon_i = \frac{k_i^2 n}{A^2}$. We denote

$$X_i = \frac{1}{a} \left(\frac{n}{A} \right)^{k_i-1} \frac{R_{k+1}^\lambda}{n^{k_i}},$$

$$Y_i = \frac{1}{a} \left(\frac{n}{A} \right)^{k_i-1} \sum_j (\alpha_j^{k_i} - (-\beta_j)^{k_i}).$$

Let us fix $a > 2$. Clearly, $|Y_i| < \frac{2}{a} < 1$ hence (28) shows that $|X_i| < 1$ if ε is smaller than some positive constant. Telescopic summation

$$X_1 \cdots X_r - Y_1 \cdots Y_r = X_1 \cdots X_{r-1} (X_r - Y_r) +$$

$$X_1 \cdots X_{r-2} (X_{r-1} - Y_{r-1}) Y_r + \cdots + (X_1 - Y_1) Y_2 \cdots Y_r$$

shows that

$$\frac{1}{a^r} \left(\frac{n}{A} \right)^{l-r} \left| \frac{\prod_{i=1}^r R_{k_i+1}^\lambda}{n^l} - \prod_{i=1}^r \left[\sum_j \alpha_j^{k_i} - \sum_j (-\beta_j)^{k_i} \right] \right| \leq \frac{2C_0(\epsilon_1 + \dots + \epsilon_r)}{a}.$$

□

5.4. Disjoint cumulants of conjugacy classes. Since this section is not central to the this paper we will be quite condensed with presenting the necessary definitions. For more details we refer to [Śni06b].

For any permutation $\pi \in S_l$ we consider the normalized conjugacy class $\Sigma_\pi \in S_n$. For normalized conjugacy classes $\Sigma_{\pi_1}, \dots, \Sigma_{\pi_r}$ we consider their *disjoint cumulant* $k^\bullet(\Sigma_{\pi_1}, \dots, \Sigma_{\pi_r})$ with respect to some given representation ρ of S_n .

One of the main results of [Śni06b] was asymptotics of the disjoint cumulant $k^\bullet(\Sigma_{\pi_1}, \dots, \Sigma_{\pi_r})$ in the case when $\pi_1, \dots, \pi_r$ are fixed and the Young diagram λ is balanced. The following Proposition (after applying similar estimates as in the proof of Theorem 1) gives a uniform estimate for such disjoint cumulants.

Proposition 22. *Let $\pi_1 \in S_{l_1}, \dots, \pi_r \in S_{l_r}$. If $\rho = \rho^\lambda$ is an irreducible representation corresponding to the Young diagram λ then*

$$k^\bullet(\Sigma_{\pi_1}, \dots, \Sigma_{\pi_r}) = \sum_{\substack{\sigma_1, \sigma_2 \in S_{l_1+\dots+l_r}, \\ \sigma_1 \sigma_2 = \pi_1 \times \dots \times \pi_r, \\ \langle \sigma_1, \sigma_2, S_{l_1} \times \dots \times S_{l_r} \rangle \text{ is transitive}}} (-1)^{|\sigma_1|} N^\lambda(\sigma_1, \sigma_2),$$

where the sum runs over all factorizations $\sigma_1 \sigma_2 = \pi_1 \dots \pi_r \in S_{l_1+\dots+l_r}$ which are transitive in a sense that the permutations σ_1, σ_2 together with $S_{l_1} \times \dots \times S_{l_r}$ act transitively on $\{1, \dots, l_1 + \dots + l_r\}$.

Proof. It is an immediate consequence of Theorem 2 and the definition of the disjoint cumulants. □

ACKNOWLEDGMENTS

Research of PŚ is supported by the MNiSW research grant P03A 013 30, by the EU Research Training Network “QP-Applications”, contract HPRN-CT-2002-00279 and by the EC Marie Curie Host Fellowship for the Transfer of Knowledge “Harmonic Analysis, Nonlinear Analysis and Probability”, contract MTKD-CT-2004-013389.

VF has been supported by the Project Polonium, one of Egide’s PHC to visit PŚ. He also would like to thank his advisor Philippe Biane.

REFERENCES

- [Bia97] Philippe Biane. Some properties of crossings and partitions. *Discrete Math.*, 175(1-3):41–53, 1997.
- [Bia98] Philippe Biane. Representations of symmetric groups and free probability. *Adv. Math.*, 138(1):126–181, 1998.
- [Bia03] Philippe Biane. Characters of symmetric groups and free cumulants. In *Asymptotic combinatorics with applications to mathematical physics (St. Petersburg, 2001)*, volume 1815 of *Lecture Notes in Math.*, pages 185–200. Springer, Berlin, 2003.
- [DH04] Ken Dykema and Uffe Haagerup. Invariant subspaces of the quasinilpotent DT-operator. *J. Funct. Anal.*, 209(2):332–366, 2004.
- [Fér06] Valentin Féray. Proof of Stanley’s conjecture about irreducible character values of the symmetric group. Preprint arXiv:math.CO/0612090, 2006.
- [GJ92] I. P. Goulden and D. M. Jackson. The combinatorial relationship between trees, cacti and certain connection coefficients for the symmetric group. *European J. Combin.*, 13(5):357–365, 1992.
- [GR05] I.P. Goulden and A. Rattan. An explicit form for Kerov’s character polynomials. Preprint math.CO/0505317, 2005.
- [Ker93] S. Kerov. The asymptotics of interlacing sequences and the growth of continual Young diagrams. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)*, 205(Differentsialnaya Geom. Gruppy Li i Mekh. 13):21–29, 179, 1993.
- [Ker99] S. Kerov. A differential model for the growth of Young diagrams. In *Proceedings of the St. Petersburg Mathematical Society, Vol. IV*, volume 188 of *Amer. Math. Soc. Transl. Ser. 2*, pages 111–130, Providence, RI, 1999. Amer. Math. Soc.
- [MR06] Cristopher Moore and Alexander Russell. On the impossibility of a quantum sieve algorithm for graph isomorphism. Preprint quant-ph/0609138, 2006.
- [Roi96] Yuval Roichman. Upper bound on the characters of the symmetric groups. *Invent. Math.*, 125(3):451–485, 1996.
- [RŚ06] Amarpreet Rattan and Piotr Śniady. Upper bound on the characters of the symmetric groups for balanced Young diagrams and a generalized Frobenius formula. Preprint math.RT/0610540, 2006.
- [Shl96] Dimitri Shlyakhtenko. Random Gaussian band matrices and freeness with amalgamation. *Internat. Math. Res. Notices*, (20):1013–1025, 1996.
- [Śni03] Piotr Śniady. Multinomial identities arising from free probability theory. *J. Combin. Theory Ser. A*, 101(1):1–19, 2003.
- [Śni06a] Piotr Śniady. Asymptotics of characters of symmetric groups, genus expansion and free probability. *Discrete Math.*, 306(7):624–665, 2006.
- [Śni06b] Piotr Śniady. Gaussian fluctuations of characters of symmetric groups and of Young diagrams. *Probab. Theory Related Fields*, 136(2):263–297, 2006.
- [Spe98] Roland Speicher. Combinatorial theory of the free product with amalgamation and operator-valued free probability theory. *Mem. Amer. Math. Soc.*, 132(627):x+88, 1998.
- [SS98] Ya. G. Sinai and A. B. Soshnikov. A refinement of Wigner’s semicircle law in a neighborhood of the spectrum edge for random symmetric matrices. *Funktsional. Anal. i Prilozhen.*, 32(2):56–79, 96, 1998.

- [Sta04] Richard P. Stanley. Irreducible symmetric group characters of rectangular shape. *Sém. Lothar. Combin.*, 50:Art. B50d, 11 pp. (electronic), 2003/04.
- [Sta06] Richard P. Stanley. A conjectured combinatorial interpretation of the normalized irreducible character values of the symmetric group. Preprint arXiv:math.CO/0606467, 2006.
- [VDN92] D. V. Voiculescu, K. J. Dykema, and A. Nica. *Free random variables*, volume 1 of *CRM Monograph Series*. American Mathematical Society, Providence, RI, 1992. A noncommutative probability approach to free products with applications to random matrices, operator algebras and harmonic analysis on free groups.
- [VK81] A. M. Vershik and S. V. Kerov. Characters and factor representations of the infinite symmetric group. *Dokl. Akad. Nauk SSSR*, 257(5):1037–1040, 1981.
- [Zvo97] A. Zvonkin. Matrix integrals and map enumeration: an accessible introduction. *Math. Comput. Modelling*, 26(8-10):281–304, 1997. Combinatorics and physics (Marseilles, 1995).

THE GASPARD–MONGE INSTITUT OF ELECTRONIQUE AND COMPUTER SCIENCE,
UNIVERSITY OF MARNE-LA-VALLÉE PARIS-EST, 77454 MARNE-LA-VALLÉE CEDEX
2, FRANCE

E-mail address: feray@ens.fr

INSTITUTE OF MATHEMATICS, UNIVERSITY OF WROCLAW, PL. GRUNWALDZKI 2/4,
50-384 WROCLAW, POLAND

E-mail address: Piotr.Sniady@math.uni.wroc.pl